

Recombinant Human KGF

Catalog No.: RP0011

Basic Information

Information


Source	<i>E.coli</i>
Description	Recombinant Human Fibroblast Growth Factor 7/Keratinocyte Growth Factor is produced by our E.coli expression system and the target gene encoding Cys32-Thr194 is expressed.
Accession	P21781
Known As	Fibroblast growth factor 7; FGF-7; Heparin-binding growth factor 7; HBGF-7; Keratinocyte growth factor; FGF7; KGF
Predicted Mol Mass	18.9 KDa
Apparent Mol Mass	17 KDa, reducing conditions

Properties


Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM Tris, 1mM EDTA, 5% Trehalose, 0.02% Tween 80, pH 8.0.
Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Endotoxin	< 1 EU/µg as determined by LAL test.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.

Experimental Data

Purity-SDS-PAGE



Bioactivity-Cell Based Assay



Greater than 95% as determined by reducing SDS-PAGE. (QC verified)

Immobilized Human KGF at 5 $\mu\text{g/mL}$ (100 $\mu\text{L/well}$) can bind Human FGFR2 beta IIIb (C-Fc). The EC50 of Human FGFR2 beta IIIb (C-Fc) is 83 ng/mL. (Regularly tested)

Background

Fibroblast growth factor 7 (FGF7) is a secreted protein which is mainly located in epithelial cells and belongs to the heparin-binding growth factors family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. FGF7 is a potent epithelial cell-specific growth factor, whose mitogenic activity is predominantly exhibited in keratinocytes but not in fibroblasts and endothelial cells. It is possible major paracrine effector of normal epithelial cell proliferation.