

Recombinant Human IL-7 (C-6His)

Catalog No.: RP0022

Basic Information

Information


Source	<i>Human Cells</i>
Description	Recombinant Human Interleukin-7 is produced by our Mammalian expression system and the target gene encoding Asp26-His177 is expressed with a 6His tag at the C-terminus.
Accession	P13232
Known As	Interleukin-7; IL-7; IL7
Predicted Mol Mass	18.4 KDa
Apparent Mol Mass	19-30 KDa, reducing conditions

Properties


Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Endotoxin	< 0.01 EU/µg as determined by LAL test.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.

Experimental Data

Purity-SDS-PAGE



Bioactivity-Cell Based Assay



Greater than 95% as determined by reducing SDS-PAGE. (QC verified) Measured in a cell proliferation assay using PHA-activated human peripheral blood lymphocytes (PBL). The ED50 for this effect is 50-300 pg/ml.

Background

Human Interleukin 7 (IL-7) is a potent lymphoid cell growth factor stimulating the proliferation of lymphoid progenitors. IL7 can associate with the hepatocyte growth factor (HGF) to form a hybrid cytokine that functions as a pre-pro-B cell growth-stimulating factor. Human IL7 cDNA encodes a 177 amino acid precursor protein containing a 25 amino acid signal peptide and a 152 amino acid mature protein. Human and mouse IL7 share 65% sequence identity in the mature region and both exhibit cross-species activity. IL-7 signals via IL-7 receptor (IL7R) activating multiple pathways including JaK/STAT and PI3K/AKT, which regulate lymphocyte survival, glucose uptake, proliferation, and differentiation. IL-7 is also associated with cytoplasmic IL2-R gamma for signal transduction.